

[← Back to Search](#)

Report: Indicators of Women in Abortion Situation

Filters: [+](#)[Issue Report](#)PERINATAL
INFORMATION
SYSTEM09/03/2023 10:44:53
v.2.4.1
http://localhost:9000/db [sip-demo]
Patients: 208884 - Pregnancies: 216132 - Newborns and terminations: 171845Latin American Centre for Perinatology
Women and Reproductive Health

Other Country

Indicators of Women in Abortion Situation

Women in abortion situation

Indicates the number of women in a legal or incomplete abortion situation

	TOTAL NUMBER CASES	% OF ALL CASES	% OF VALID CASES
# Abortions	23510	96.30	
no data	892	3.65	
		n cases = 24413	

Number of women who had a uterine evacuation

Indicates the number of women who had a uterine evacuation

	TOTAL NUMBER CASES	% OF ALL CASES	% OF VALID CASES
# Uterine evacuation	20798	88.46	90.34
no data	489	2.08	
		n cases = 23510	n valid cases = 23021

Indicators of maternal morbidity and mortality

Maternal mortality ratio at discharge

The frequency of occurrence of deaths in women in abortion situation

	TOTAL NUMBER CASES	% OF ALL CASES	% OF VALID CASES
# Dead woman at discharge	7	0.03	0.07
no data	12966	55.15	
		n cases = 23510	n valid cases = 10371

Abortion-related morbidity

The frequency of women presenting complications related to abortion in women in abortion situation

	TOTAL NUMBER CASES	% OF ALL CASES	% OF VALID CASES
# Complications	696	2.96	2.98
no data	153	0.65	
		n cases = 23510	n valid cases = 23357

Severe abortion-related morbidity (Near Miss)

The frequency of women presenting severe complications (Near Miss) related to abortion in women in abortion situation

	TOTAL NUMBER CASES	% OF ALL CASES	% OF VALID CASES
# Near Miss	151	0.64	0.75
no data	3243	13.79	
		n cases = 23510	n valid cases = 20267

Preventable and non-preventable adverse effects associated with performing uterine evacuation

The frequency of women experiencing adverse effects in women who had a uterine evacuation

	TOTAL NUMBER CASES	% OF ALL CASES	% OF VALID CASES
# Adverse effect	110	0.53	0.54
no data	392	1.88	
		n cases = 20798	n valid cases = 20406

Quality pre and post abortion clinical practices

Types of procedures performed

Frequency of types of procedures performed in women who underwent a uterine evacuation

	TOTAL NUMBER CASES	% OF ALL CASES
# MVA or EVA	15799	75.96
# D&C	1367	6.57
# MED	3816	18.35
# More than one method	657	3.16
		n cases = 20798

Use of pain relievers

Frequency of analgesic use in women in abortion situation

	TOTAL NUMBER CASES	% OF ALL CASES	% OF VALID CASES
# Analgesia performed	20281	86.27	87.04
# Analgesia performed with no data	208	0.88	
		n cases = 23510	n valid cases = 23302

Use of prophylactic antibiotics

Frequency of use of prophylactic antibiotics in women who underwent MVA or EVA or D&C

	TOTAL NUMBER CASES	% OF ALL CASES	% OF VALID CASES
# Antibiotics	4960	21.10	34.30
# Antibiotics with no data	2646	11.25	
		n cases = 17107	n valid cases = 14461

Frequency of women receiving oral or written contraceptive counseling before discharge in the total number of women in abortion situation

???

	TOTAL NUMBER CASES	% OF ALL CASES	% OF VALID CASES
# Contraception counseling	15883	67.56	71.28
# Contraception counseling with no data	458	15.41	
		n cases = 23510	n valid cases = 22283

Immediate start of a contraceptive method (CM)

Frequency of women who start a contraceptive method at discharge in women in abortion situation

	TOTAL NUMBER CASES	% OF ALL CASES	% OF VALID CASES
# CM start	13547	57.62	59.60
# CM start with no data	779	3.31	
		n cases = 23510	n valid cases = 22731

Immediate start of contraceptive method in women under 20 years of age

Frequency of women <20 years of age who start a contraceptive method at discharge in the total number of women in abortion situation

	TOTAL NUMBER CASES	% OF ALL CASES	% OF VALID CASES
# CM start < 20 years	1502	50.54	52.33
# CM start < 20 years with no data	15	0.50	
		n cases = 2972	n valid cases = 2870