

Back to Search

Report: Dominican Republic - Attention quality

Filters: +

Issue Report



PERINATAL INFORMATION SYSTEM

08/03/2023 21:39:29 v2.4.1 http://localhost:9000/db/isp-demo

Patients: 208884 - Pregnancies: 216132 - Newborns and Terminations: 171845

Latin American Centre for Perinatology Women and Reproductive Health

Other Country

Dominican Republic Attention quality

Failure of contraceptive method in pregnant women who did not plan their pregnancy

Table with 4 columns: Contraceptive method, Total number cases, % of all cases, % of valid cases. Rows include Barrier, IUD, Hormone, Emergency, Natural, and no data.

GA at first visit

Table with 4 columns: Gestational age, Total number cases, % of all cases, % of valid cases. Rows include <= 13, 14 to 27, 28 or more, and no data.

Syphilis Screening and Treatment.

Total of the base. Pregnant women who have had at least one diagnostic test during pregnancy

Table with 4 columns: Diagnostic test result, Total number cases, Coverage of diagnostic tests, % of total cases with diagnostic test. Rows include Positive, Negative, and Cases with diagnostic test.

Test Reactive

Table with 4 columns: Pregnant treatment, Total number cases, % of all cases, % of valid cases. Rows include Yes, No, Not known, Not applicable, and no data.

VDRL of newborn. In pregnant women who have had at least one positive test during pregnancy

Table with 4 columns: Neonatal VDRL, Total number cases, % of all cases, % of valid cases. Rows include Positive, Negative, Not done, and no data.

In total base. Infants with a diagnostic test at birth in cord blood

Table with 4 columns: Newborn's VDRL, Total number cases, VDRL coverage, % of total number of cases with VDRL. Rows include Neonatal screening - positive, Negative, and Total number of cases.

HIV

HIV test performed

Table with 4 columns: HIV test status, Total number cases, % of all cases, % of valid cases. Rows include Requested, Not requested, Not applicable, and no data.

HIV + during pregnancy

Table with 4 columns: HIV status, Total number cases, Coverage of diagnostic tests, % of total cases with diagnostic test. Rows include Personal history HIV+, Positive, Negative, and Cases with diagnostic test.

Maternal Pathologies in Pregnancy and Childbirth

Table with 5 columns: Pathology, N, %, No data, % no data. Rows include With some pathology, Multiple pregnancy, Previous Hypertension, Preeclampsia, Eclampsia, Heart disease, Diabetes, Urinary infection, Fetal growth restriction, Threatened preterm delivery, Hemorrhage, Premature ruptured membranes, Puerperal infection, Postpartum Hemorrhage, and Other diseases.

Magnesium sulfate in eclampsia

Table with 4 columns: Magnesium sulfate use, Total number cases, % of all cases, % of valid cases. Rows include Yes, No, and no data.

Magnesium sulfate in preeclampsia

Table with 4 columns: Magnesium sulfate use, Total number cases, % of all cases, % of valid cases. Rows include Yes, No, and no data.

Hemorrhage: use of oxytocics

Table with 4 columns: Oxytocic use, Total number cases, % of all cases, % of valid cases. Rows include Uterotonic for tto. hemorrhage = no, Uterotonic for tto. hemorrhage = yes, and no data.

Infection / puerperal sepsis: use of parenteral antibiotics

Table with 4 columns: Antibiotic use, Total number cases, % of all cases, % of valid cases. Rows include Use of ATB IV for tto. of infectious complication = no, Use of ATB IV for tto. of infectious complication = yes, and no data.

Prevention of postpartum hemorrhage. MATEP: use of oxytocin

Table with 4 columns: Oxytocin use, Total number cases, % of all cases, % of valid cases. Rows include Oxytocics after expulsion of placenta = no, Oxytocics after expulsion of placenta = yes, and no data.

Early cord clamping

Table with 4 columns: Cord clamping status, Total number cases, % of all cases, % of valid cases. Rows include Yes, No, and no data.

Maternal Mortality

Causes of Maternal Mortality

Table with 4 columns: Cause of mortality, Total number cases, % of all cases, % of valid cases. Rows include Hypertensive disorders, Infections, 1st trimester hemorrhage, 2nd trimester hemorrhage, 3rd trimester hemorrhage, Obstetric complications, Metabolic disorders, and Other disorders.

Reason for Extreme Maternal Morbidity (Near Miss)

Table with 4 columns: Reason for morbidity, Total number cases, % of all cases, % of valid cases. Rows include Variables to Identify Near Miss cases = yes.

Extreme Maternal Morbidity (Near Miss) for cause

Table with 4 columns: Cause of morbidity, Total number cases, % of all cases, % of valid cases. Rows include Hypertensive disorders, Infections, 1st trimester hemorrhage, 2nd trimester hemorrhage, 3rd trimester hemorrhage, Obstetric complications, Metabolic disorders, and Other disorders.

Extreme Maternal Morbidity (Near Miss) by age group

Table with 4 columns: Age group, Total number cases, % of all cases, % of valid cases. Rows include <15 years, 15-19 years, 20-34 years, 35-44 years, 45-49 years, and >49 years.

Extreme Maternal Morbidity Ratio (Near Miss) / Maternal Death

Table with 4 columns: Ratio, Total number cases, % of all cases, % of valid cases. Rows include Variables to Identify Near Miss cases = yes.

Lethality Rates

Pre-Eclampsia Lethality Rate

Table with 4 columns: Lethality status, Total number cases, % of all cases, % of valid cases. Rows include Yes, No, and no data.

Eclampsia Lethality Rates

Table with 4 columns: Lethality status, Total number cases, % of all cases, % of valid cases. Rows include Yes, No, and no data.

Postpartum Infection Lethality Rate

Table with 4 columns: Lethality status, Total number cases, % of all cases, % of valid cases. Rows include Yes, No, and no data.

Postpartum Hemorrhage Lethality Rate

Table with 4 columns: Lethality status, Total number cases, % of all cases, % of valid cases. Rows include Yes, No, and no data.

Termination

Table with 4 columns: Termination method, Total number cases, % of all cases, % of valid cases. Rows include Spontaneous, C-Section, Forceps, Vacuum, Other, and no data.

Mortality

Table with 4 columns: Mortality type, N, Rate per thousand live births + deaths, % of all cases. Rows include Fetal Mortality, Early Neonatal Mortality, Perinatal Mortality, Late neonatal mortality rate, Early Neonatal Mortality, and Maternal deaths.

Antenatal corticosteroids

Table with 4 columns: Corticosteroid use, Total number cases, % of all cases, % of valid cases. Rows include Yes, No, and no data.

Gestational Age

Table with 4 columns: Gestational age, Total number cases, % of all cases, % of valid cases. Rows include < 22 weeks, 22-27 weeks, 28-36 weeks, 37-41 weeks, > 41 weeks, and no data.

Birth Weight

Table with 4 columns: Birth weight, Total number cases, % of all cases, % of valid cases. Rows include < 500, 500 - 999, 1000 - 1499, 1500 - 1999, 2000 - 2499, 2500 - 3999, >= 4000, and no data.

Apgar Score

Apgar Score 1st minute. Livebirths only.

Table with 4 columns: Apgar score, Total number cases, % of all cases, % of valid cases. Rows include Present, 0 - 3, 4 - 6, 7 - 10, and no data.

Apgar Score 5th minute. Livebirths only.

Table with 4 columns: Apgar score, Total number cases, % of all cases, % of valid cases. Rows include Present, 0 - 3, 4 - 6, 7 - 10, and no data.

Method of Family Planning at discharge

Focus on teenagers

Table with 5 columns: Requested method, %, Received method, % of valid cases. Rows include OCDs pill, Injection, Implant, Another hormonal therapy, Another barrier method, Condom, IUD, VSS:male, VSS:female, and Abstinence.

